

Pediatric Prescribing Recommendations

Procedure	Doses [^] of Opioid
Dental	
Dental Extraction	0
General	
Appendectomy	0
Inguinal Hernia	0
Umbilical Hernia	0
Maxillofacial	
LeFort and/or Sagittal Split Osteotomy	0 - 7 pills ^{^^}
Oral-Maxillofacial Bone Cyst Removal	0
Orthopedics	
Epiphysiodesis	0 - 1 pills ^{^^}
Hand Surgery	0 - 4
Knee Surgery - Arthroscopic, Open, Ankle Antroscopy	0 - 14 pills ^{^^}
ORIF - Elbow, Humerus, Radius, Ulna	0 - 2
ORIF - Hip	0 - 3
ORIF - Tibia, Ankle, Femur	0 - 9 pills ^{^^}
Orthopedic Bone Lesion Removal	0 - 2
Orthopedic Hardware Removal	0
Osteotomy - Rotation Femur/Tibia, Shortening Femur/Tibia	0 - 4
Percutaneous Pinning - Wrist, Elbow, Humerus	0
Orthopedic Tendon / Soft Tissue Procedure	0
Trigger Finger Release	0

Procedure	Doses [^] of Opioid
Otolaryngology	
Adenoidectomy	0
Mastoidectomy	0
Neck Surgery	0
Otoplasty	0 - 2
Tonsillectomy (11 years old and younger)	0 - 4
Tonsillectomy (12 years old and greater)	0 - 12
Tympanoplasty	0
Plastics	
Alveolar Cleft Bone Graft	0 - 5
Cleft Lip	0 - 5
Palatoplasty and Pharyngoplasty	0 - 4
Urology	
Circumcision	0
Hydrocelectomy	0
Hypospadias Repair	0
Orchiopexy	0
Penile Surgery - Chordee Release, Meatoplasty or Meatotomy	0
Testicular Torsion Repair	0 - 1
Ureteral Reimplant	0 - 2
Body	
Soft Tissue Procedure	0

[^] In patients weighing less than 50 kg, we define a dose of opioids as oxycodone ranging from 0.05 mg/kg to 0.1 mg/kg. For patients >50 kg, a dose is 5 mg of oxycodone. When patients weighing less than 50 kg fall into a BMI category >95th percentile for age, either a lower mg/kg starting dose or ideal body weight/lean body mass should be used to determine total dose. A lower mg/kg starting point should also be considered in those with comorbidities predisposing to respiratory depression.

^{^^} 5 mg Oxycodone pill